

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show 20 Sort by Send to

All: 1 Review: 0

About Entrez  
NCBI Toolbar

Text Version

Entrez PubMed

Overview  
Help | FAQ  
Tutorials  
New/Noteworthy  
E-Utilities

PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
Special Queries  
LinkOut  
My NCBI

Related Resources

Order Documents  
NLM Mobile  
NLM Catalog  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

1: Biol Psychiatry. 2001 Aug 1;50(3):205-16.

[Related Articles, Links](#)

ELSEVIER  
FULL-TEXT ARTICLE

**Dawn simulation and bright light in the treatment of SAD: a controlled study.**

**Avery DH, Eder DN, Bolte MA, Hellekson CJ, Dunner DL, Vitiello MV, Prinz PN.**

Department of Psychiatry and Behavioral Sciences, University of Washington School of Medicine, Harborview Medical Center, Seattle, Washington 98104-2499, USA.

**BACKGROUND:** Some small controlled studies have found that dawn simulation is effective in treating seasonal affective disorder (SAD). With a larger sample size and a longer duration of treatment, we compared dawn simulation with bright light therapy and a placebo condition in patients with SAD. **METHOD:** Medication-free patients with SAD were randomly assigned to one of three conditions: bright light therapy (10,000 lux for 30 min, from 6:00 AM to 6:30 AM), dawn simulation (1.5 hour dawn signal from 4:30 AM to 6:00 AM peaking at 250 lux), and a placebo condition, a dim red light (1.5 hour dawn signal from 4:30 am to 6:00 AM peaking at 0.5 lux.) Over the subsequent 6 weeks, the subjects were blindly rated by a psychiatrist using the Structured Interview Guide for the Hamilton Depression Rating-Seasonal Affective Disorder Version (SIGH-SAD). We modeled the profiles of the remissions (SIGH-SAD < or = 8) and response (> or =50% decrease in SIGH-SAD) to treatment over time using Cox proportional hazards models. **RESULTS:** The sample consisted of 95 subjects who were randomized to the three conditions: bright light (n = 33), dawn simulation (n = 31) and placebo (n = 31). Dawn simulation was associated with greater remission (p <.05) and response (p <.001) rates compared to the placebo. Bright light did not differ significantly from the placebo. Dawn simulation was associated with greater remission (p <.01) and response (p <.001) rates compared to the bright light therapy. The mean daily hours of sunshine during the week before each visit were associated with a significant increase in likelihood of both remission (p <.001) and response (p <.001). **CONCLUSIONS:** Dawn simulation was associated with greater remission and response rates compared to the placebo and compared to bright light therapy. The hours of sunshine during the week before each assessment were associated with a positive clinical response.

Publication Types:

- [Clinical Trial](#)
- [Randomized Controlled Trial](#)

PMID: 11513820 [PubMed - indexed for MEDLINE]